

literatuurlijst

Alcohol in mondspoelmiddelen

Januari 2009

samengesteld door:



The Oral Health Experts

Lichtschip 15, 3991 CP Houten
tel.: +31 (0) 30 6343154
fax.: + 31(0) 30 6343145
E-mail: info@dentaid.nl / info@dentaid.be
www.dentaid.nl

1	Arweiler N.B. et al.	Alcohol-free mouthrinse solutions to reduce supragingival plaque regrowth and vitality A controlled clinical study.	J. Clin. Periodontology	2001	28:168-174
2	Borrajó L. et al.	Efficacy of Chlorhexidine Mouthrinses with and without Alcohol; a Clinical Study.	J. Clin. Periodontology	2002	72:317-321
3	Winn D.M. et al.	Mouthwash use and oral conditions in the risk of oral and pharyngeal cancer.	Cancer Res.	1991	1;51(11):3044-7
4	Weaver A. et al.	Mouthwash and oral cancer: carcinogen or coincidence ?	J Oral Surg.	1979	37(4):250-3
5	Kabat G.C. et al.	Risk factors for oral cancer in woman.	Cancer Res.	1989	15;49(10)2803-6
6	Winn D.M. et al.	Moutwash in the etiology of oral cancer in Perto Rico.	Cancer Res.	2001	12(5):419-29
7	Weiner R.. et al.	The effect of alcoholic and nonalcoholic mouthwashes on heat-treated composite resin.	Opper Dent.	1997	22(6):249-53
8	Almerich J.M. et al.	Influence of alcohol in mouthwashes containing triclosan and zinc: an experimental gingivitis study.	J Clin Periodontology	2005	32(6):539-44
9	McCullough M.J..et al.	The role of alcohol in oral carcinogenesis with particular reference to alcohol-containing mouthwashes	Australian Dental Journal	2008	10.1111/j.1834-7819.2008.00070.x
10	Guha N. et al.	Oral Health and Risk of Squamous Cell Carcinoma of the Head and Neck and Esophagus : Results of Two Multicentric Case-Control Studies	American Journal of Epideminologie	2007	166:1159-1173